

ISSN 0960-8923, Volume 20, Number 8

OBESITY SURGERY
The Journal of Metabolic Surgery and Allied Care

Volume 20 Number 8 August 2010 ISSN 0960-8923

An International Surgical Journal for Research and Treatment of Massive Obesity

*Official Journal of the International Federation
for the Surgery of Obesity and metabolic disorders*



ABSTRACTS FROM THE IFSO XV WORLD CONGRESS, SEPTEMBER 3-7, 2010

ESOPHAGEAL PERISTALSIS AND IN VIVO INTRABAND PRESSURE MEASUREMENTS

GLUCOSE-STIMULATED INSULIN SECRETION IN GASTRIC BYPASS PATIENTS WITH HYPOGLYCEMIC SYNDROME

ASSESSMENT OF THE APPLICATION OF THE INTRAGASTRIC BALLOON TOGETHER WITH SIBUTRAMINE

HEALTH-RELATED QUALITY OF LIFE IN ADULT PATIENTS COMING FOR BARIATRIC SURGERY

IMPACT ON OBESITY-RELATED ILLNESSES AND QUALITY OF LIFE FOLLOWING INTRAGASTRIC BALLOON

NUTRITIONAL PYRAMID FOR POST-GASTRIC BYPASS PATIENTS

WHO UNDERGOES BARIATRIC SURGERY?

LAPAROSCOPIC CRURAL CLOSURE IN THE MANAGEMENT OF HIATAL HERNIA IN SLEEVE GASTRECTOMY PATIENTS

 Springer

Official Journal of the International Federation for the Surgery of Obesity and metabolic disorders, Obesity Surgery Society of Australia and New Zealand, Società Italiana di Chirurgia dell'Obesità, Österreichische Gesellschaft für Adipositaschirurgie, Sociedade Brasileira de Cirurgia Bariátrica e Metabólica, BeNeLux Association of Bariatric Surgeons, Deutsche Gesellschaft für Chirurgie der Adipositas, Czech Society for the Study of Obesity, Colegio Mexicano Cirugía de la Obesidad A.C., Sociedad Española de Cirugía de la Obesidad, Greek Society for Bariatric and Metabolic Surgery, Israeli Association for Obesity Surgery, Sociedade Paragvaya de Cirurgia de la Obesidad, Japanese Society for the Study of Obesity, Société Française de Chirurgie de l'Obésité, Yugoslav Obesity Surgery Society, Swiss Study Group for Morbid Obesity, Turkish Obesity Surgery Society, Egyptian Society for Bariatric Surgery, Society of Bariatric Surgeons of Russia, Ukrainian Association of Bariatric Surgery, Sociedad Argentina de Cirugía de la Obesidad, Kuwait Society for Bariatric Surgery, Portuguese Obesity Surgery Society, Polish Association for Prevention and Treatment of Obesity, British Obesity Surgery Society, Hungarian Bariatric Surgery Foundation, Asociación Paraguaya de Cirugía Bariátrica, Asociación Peruana de Cirugía Bariátrica y Obesidad Severa, Obesity Surgery Society of India, Departamento Cirugía Bariátrica Chile, Romanian Association of Bariatric Surgery, Asia Pacific Bariatric Surgical Society, Sección de Cirugía Bariátrica de la Sociedad Venezolana de Cirugía, South African Association for Obesity and Metabolism, The International Society for the Perioperative Care of the Obese Patient

**This article was published in the above mentioned Springer issue.
The material, including all portions thereof, is protected by copyright;
all rights are held exclusively by Springer Science + Business Media.
The material is for personal use only;
commercial use is not permitted.
Unauthorized reproduction, transfer and/or use
may be a violation of criminal as well as civil law.**

Laparoscopic Adjustable Banded Sleeve Gastrectomy as a Primary Procedure for the Super-Super Obese (Body Mass Index >60 kg/m²)

Sanjay Agrawal · Els Van Dessel · Faki Akin ·
Sebastiaan Van Cauwenberge · Bruno Dillemans

Published online: 14 May 2010
© Springer Science+Business Media, LLC 2010

Abstract Isolated laparoscopic sleeve gastrectomy is increasingly being used for the treatment of morbid obesity. However, doubts still persist regarding long-term weight loss, and the 5-year results are awaited. Whether the aetiology of failed excess weight loss is the result of an inadequate sleeve or attributable to dilatation of the sleeve is not clear. In an effort to prevent gastric dilatation and increase gastric restriction to promote further weight loss in the long term, we performed a combined procedure of laparoscopic adjustable gastric banding with sleeve gastrectomy. The patient was a 39-year-old woman with a life-long history of obesity and a body mass index of 79.8 kg/m². The surgical technique of the laparoscopic adjustable gastric banded sleeve gastrectomy is described. There were no immediate complications, and the patient was discharged home on the third postoperative day. She is doing extremely well on clinic follow-up at 6 weeks. To the best of our knowledge, laparoscopic adjustable gastric banded sleeve gastrectomy, as a primary operation, has not been described in the literature. It is hoped that this combined procedure will be most useful in the super-super obese (body mass

index >60) patients. More patients with a long-term follow-up are necessary to provide definitive conclusions regarding long-term benefits and complications of this combined bariatric procedure.

Keywords Adjustable gastric band · Sleeve gastrectomy · Morbid obesity · Super-super obesity · Laparoscopic

Introduction

Sleeve gastrectomy (SG) was first described as a modification to the bilio-pancreatic diversion (BPD) and combined with a duodenal switch (DS) in 1998 [1, 2]. It was first performed laparoscopically in July 1999 and reported the following year [3]. Laparoscopic sleeve gastrectomy (LSG), as a primary operation in the management of morbid obesity, was first reported in 2003 [4]. Since then, multiple recent reports have documented SG as single therapy in the treatment of morbid obesity [5, 6]. With increasing experience, a number of complications have been reported with SG including dilatation of the remaining stomach [7–9]. Also, doubts still persist regarding long-term weight loss, and the 5-year results are awaited. Whether the aetiology of failed excess weight loss is the result of an inadequate sleeve or attributable to dilatation or hypertrophy of the sleeve is not clear. In an effort to prevent gastric dilatation and increase gastric restriction to promote weight loss in the long term, we performed a combined procedure of laparoscopic adjustable gastric banding with sleeve gastrectomy as described below.

The patient was a 39-year-old woman with a life-long history of obesity and a body mass index of 79.8 kg/m² (241 kg with 174 cm height). Her co-morbidities included asthma, musculo-skeletal complaints, and depression on

Financial support None.

Submitted to the meeting of the 4th European IFSO Congress, 22–24 April, 2010, Lausanne, Switzerland.

S. Agrawal
Department of Bariatric Surgery, Homerton University Hospital,
London, UK
e-mail: sanju_agrawal@hotmail.com

E. Van Dessel · F. Akin · S. Van Cauwenberge · B. Dillemans (✉)
Department of General Surgery, AZ Sint-Jan Hospital AV,
Bruges, Belgium
e-mail: bruno.dillemans@azbrugge.be

treatment. Following a thorough informed discussion with her regarding options of surgical treatment and subsequent multidisciplinary meeting, it was decided to perform either a laparoscopic long-limb Roux-en-Y gastric bypass or laparoscopic sleeve gastrectomy with an adjustable band as a primary procedure.

Surgical Technique

The patient was placed in the supine position, split-leg with reverse Trendelenberg position along with slight flexion of the hip to help increase surgical abdominal workspace [10]. The surgeon [B.D.] stood between the legs. A video monitor was positioned at the level of the patient's head. A 30° angled scope was used. Abdominal insufflation with carbon dioxide was achieved using a Veress needle. Intra-abdominal pressure was maintained at 15 mm Hg. A five-port technique was employed: a 10-mm port 10 cm below the xiphoid process, a 5-mm port high in the epigastrium on the midline, a 12-mm port in the right upper quadrant, and two 12-mm ports in the left upper quadrant. All the three 12-mm ports were placed on the same line of the 10-mm port.

Intraoperatively, extensive adhesions in the lower abdomen due to previous surgery prevented us from continuing with Roux-en-Y gastric bypass operation. The SG was done from approximately 5 cm proximal to the pylorus to the gastroesophageal junction (angle of His) over a 34-Fr bougie using a linear stapler (Echelon 60 Endopath stapler, Ethicon; Cincinnati, OH, USA). An adjustable gastric band (AGB) (HAGE, Helioscopie, France) was placed through a small opening between the blood vessels on the lesser curve and the sleeve approximately 6 cm below the gastroesophageal junction (Figs. 1 and 2). The band was sutured laterally with a couple of nonabsorbable sutures (2/0

Fig. 1 Laparoscopic adjustable gastric banded sleeve gastrectomy



Fig. 2 Laparoscopic adjustable gastric banded sleeve gastrectomy

ethibond, Ethicon) to the peripancreatic tissues. Test for leakage was done by forcefully injecting methylene blue through the orogastric tube after a clamp was placed to temporarily occlude the pylorus to achieve complete distension of the remaining stomach. The subcutaneous reservoir port was secured to the anterior rectus sheath in a midclavicular plane, and hollow tubing was used to connect the band to the port. The band was not filled at operation. Postoperatively, she was allowed free fluids from the second postoperative day and was discharged home on the third postoperative day. She was advised to be on liquid diet for 2 weeks and then advance to a pureed and solid diet over the next 4 weeks. She is doing extremely well on clinic follow-up at 6 weeks.

Discussion

To the best of our knowledge, a combination of an AGB with LSG, as a primary operation for the treatment of morbid obesity, has not been described in the literature. The placement of an adjustable band around the upper sleeve will further limit the volume of food intake and prevent dilatation or hypertrophy of the gastric sleeve distal to the band in the long term. With an LAGB component, there is a sequential action mechanism: LSG is mainly active at the beginning, while adjustable gastric restriction from an AGB will result in further weight loss following this period. Also, the adjustable band will help for weight maintenance, especially when restriction starts to fail and weight regain occurs due to gastric dilatation. The procedure thus combines the potential benefits of SG and an AGB.

Likewise, as a primary operation, Gagner et al. [11] as well as Vassallo et al. [12] reported morbidly obese patients who underwent a LAGB with BPD and DS and had satisfactory weight loss. Also, Gabriel et al. [13]

performed a LAGB with a BPD and reported as much as 99.6% reduction in excess weight at 2 years. Combining previous gastric restrictive surgery with a subsequent BPD has also been previously described [14, 15]. These procedures are more invasive than this technique but represent other means for a combined restrictive/malabsorptive procedure.

In an effort to prevent gastric dilatation, several modifications of the SG can be done, thereby improving long-term results. There has been a recent report of using human collagen as a band for sleeve gastrectomy at the primary operation [16]. Greenstein et al. [17] recently reported placement of a laparoscopically inserted gastric band around the upper part of the stomach that had a previous sleeve gastrectomy that had become dilated and was associated with poor weight loss. At 9 months, the patient had a 57% loss of excess body weight, suggesting the utility of combining banding with sleeve gastrectomy. Another modification of a band with a sleeve gastrectomy was evaluated by de Paula et al. [18] who reported a series of 19 patients who had a laparoscopic sleeve gastrectomy over a 30-Fr orogastric calibration tube with placement of a silicone band around the stomach 3–4 cm below the cardia. However, they also interposed a 100-cm long segment of ileum into the jejunum 50 cm from the ligament of Treitz to provide a 'neuroendocrine brake'. Other techniques to prevent gastric dilatation include wrapping the gastric sleeve in polytetrafluoroethylene dual mesh. A preliminary study from New York found significantly less weight gain in a porcine model after 8 weeks [19].

The AGB, however, has been associated with late complications, including slippage and erosion of the band. Since the band is placed through a small opening between the blood vessels immediately adjacent to the stomach and the lesser curve and fixed laterally with the peripancreatic tissues, the chance of slippage is expected to be low. Whether or not late complications will occur remains to be seen.

Conclusion

The insertion of an AGB is feasible and safe during a LSG at the primary operation, with no immediate major complications in this case. We assume that combining an adjustable band can reduce the incidence of failed sleeve gastrectomies, leading to better excess weight loss results than those of a sleeve alone. It is hoped that this combined procedure will be most useful in the super-super obese (body mass index > 60) patients. More patients with a long-term follow-up are necessary to provide definitive conclusions regarding long-term benefits and complications of this combined bariatric procedure.

Acknowledgement S.A. was supported by a travelling fellowship grant from Covidien plc, United Kingdom (UK) and awarded by the Association of Surgeons in Training (ASiT), UK.

Conflict of interest disclosure The authors declare that they have no conflict of interest.

References

- Hess DS, Hess DW. Biliopancreatic diversion with a duodenal switch. *Obes Surg.* 1998;8:267–82.
- Marceau P, Hould FS, Simard S, et al. Biliopancreatic diversion with duodenal switch. *World J Surg.* 1998;22:947–54.
- Ren CJ, Patterson E, Gagner M. Early results of laparoscopic biliopancreatic diversion with duodenal switch: a case series of 40 consecutive patients. *Obes Surg.* 2000;10:514–23. discussion 524.
- Regan JP, Inabnet WB, Gagner M, et al. Early experience with two-stage laparoscopic Roux-en-Y gastric bypass as an alternative in the super-super obese patient. *Obes Surg.* 2003;13(6):861–4.
- Baltasar A, Serra C, Perez N, et al. Laparoscopic sleeve gastrectomy: a multipurpose bariatric operation. *Obes Surg.* 2005;15:1124–8.
- Hamoui N, Anthonie GJ, Kaufman HS, et al. Sleeve gastrectomy in the high-risk patient. *Obes Surg.* 2006;16:1445–9.
- Langer FB, Bohdjalian A, Felberbauer FX, et al. Does gastric dilatation limit the success of sleeve gastrectomy as a sole operation for morbid obesity? *Obes Surg.* 2006;16:166–71.
- Baltasar A, Serra C, Perez N, et al. Re-sleeve gastrectomy. *Obes Surg.* 2006;16:1535–8.
- Nocca D, Krawczykowsky D, Bomans B, et al. A prospective multicenter study of 163 sleeve gastrectomies: results at 1 and 2 years. *Obes Surg.* 2008;18:560–5.
- Mulier JP, Dillemans B, Luyten A, et al. Horizontal positioning of the trunk and maximal leg flexion increases the laparoscopic workspace for bariatric surgery. *Obes Surg.* 2008;18:442.
- Gagner M, Steffen R, Biertho L, et al. Laparoscopic adjustable gastric banding with duodenal switch for morbid obesity: technique and preliminary results. *Obes Surg.* 2004;13:444–9.
- Vassallo C, Negri L, Rovatti P, et al. Biliopancreatic diversion with transitory gastric restriction and duodenal bulb preservation: 88 patients since 1992. *Obes Surg.* 2004;14:773–6.
- Gabriel SG, Karaindros CA, Papaioannou MA, et al. Biliopancreatic diversion with duodenal switch combined with laparoscopic adjustable gastric banding. *Obes Surg.* 2005;15:517–22.
- Yashkov YI, Oppel TA, Shishlo LA, et al. Improvement of weight loss and metabolic effects of vertical banded gastroplasty by an added duodenal switch procedure. *Obes Surg.* 2001;11:635–9.
- Slater GH, Fielding G. Combining laparoscopic adjustable gastric banding and biliopancreatic diversion after failed bariatric surgery. *Obes Surg.* 2004;14:677–82.
- Alexander JW, Martin Hawver LR, Goodman HR. Banded sleeve gastrectomy—initial experience. *Obes Surg.* 2009;19:1591–6.
- Greenstein AJ, Vine AJ, Jacob BP. When sleeve gastrectomy fails: adding a laparoscopic adjustable gastric band to increase restriction. *Surg Endosc.* 2009;23:884.
- de Paula AL, Macedo ALV, Prudente AS, et al. Laparoscopic sleeve gastrectomy with ileal interposition ('neuroendocrine brake')—pilot study of a new operation. *Surg Obes Relat Dis.* 2006;2:464–7.
- Ueda K, Gagner M, Milone L, et al. Sleeve gastrectomy with wrapping using polytetrafluoroethylene to prevent gastric enlargement in a porcine model. *Surg Obes Relat Dis.* 2008;4(2):84–90.